

# Refined Razumov–Stroganov conjectures for open boundaries

Jan de Gier<sup>1</sup> and Vladimir Rittenberg<sup>2</sup>

<sup>1</sup> Department of Mathematics and Statistics, The University of Melbourne, VIC 3010, Australia

<sup>2</sup> Physikalisches Institut, Universität Bonn, Nussallee 12, Bonn, Germany  
E-mail: [degier@ms.unimelb.edu.au](mailto:degier@ms.unimelb.edu.au) and [vladimir@th.physik.uni-bonn.de](mailto:vladimir@th.physik.uni-bonn.de)

Received 25 August 2004

Accepted 21 September 2004

Published 29 September 2004

Online at [stacks.iop.org/JSTAT/2004/P09009](http://stacks.iop.org/JSTAT/2004/P09009)

doi:10.1088/1742-5468/2004/09/P09009

**Abstract.** Recently it has been conjectured that the ground state of a Markovian Hamiltonian, with one boundary operator, acting in a link pattern space is related to vertically and horizontally symmetric alternating-sign matrices (equivalently fully packed loop configurations (FPL) on a grid with special boundaries). We extend this conjecture by introducing an arbitrary boundary parameter. We show that the parameter dependent ground state is related to refined vertically symmetric alternating-sign matrices, i.e. with prescribed configurations (respectively, prescribed FPL configurations) in the next to central row.

We also conjecture a relation between the ground state of a Markovian Hamiltonian with two boundary operators and arbitrary coefficients and some doubly refined (dependence on two parameters) FPL configurations. Our conjectures might be useful in the study of ground states of the O(1) and XXZ models, as well as the stationary states of Raise and Peel models.

**Keywords:** algebraic structures of integrable models, loop models and polymers, quantum integrability (Bethe ansatz), solvable lattice models

**ArXiv ePrint:** [math-ph/0408042](http://arxiv.org/abs/math-ph/0408042)

---

**Contents**

<b>1. Introduction</b>	<b>2</b>
<b>2. The refined one-boundary RS conjecture</b>	<b>3</b>
2.1. Fully packed loops . . . . .	5
2.2. A refined conjecture . . . . .	7
2.3. An application . . . . .	8
2.4. The relation to the XXZ chain with diagonal boundary conditions . . . . .	9
<b>3. The two-boundary case</b>	<b>10</b>
<b>4. Conclusion</b>	<b>12</b>
<b>Acknowledgments</b>	<b>13</b>
<b>Appendix. The eigenvector <math>\psi_0</math></b>	<b>13</b>
<b>References</b>	<b>13</b>

---

**1. Introduction**

In a remarkable paper, Razumov and Stroganov (RS) [1] have looked at the ground state wavefunction of the ferromagnetic one-dimensional XXZ spin 1/2 chain with the asymmetry parameter  $\Delta = -1/2$ , odd number of sites and periodic boundary condition. They noticed that for a small number of lattice sites  $L = 2n + 1$  (this was all numerics), the largest component and the normalization are related to the number of  $n \times n$  alternating-sign matrices (ASMs) [2]. They conjectured that these coincidences are valid for any number of sites. This conjecture, though not yet proven, triggered a lot of other conjectures which brought together the study of quantum chains and combinatorics.

It was realized in [3] that similar conjectures can be made in the  $O(1)$  loop model with various boundary conditions. This led RS [4], using the bijection between ASMs and fully packed loops (FPLs) on a grid with special boundary conditions, to a new conjecture (called hereafter the RS conjecture) which gives a much deeper connection between the ground state wavefunction and ASMs. Conjectures similar to the one of RS were also made for Hamiltonians with different boundary conditions [5, 6] relating them to different symmetry classes of ASMs.

In another development (relevant to the present paper), it was understood that the results obtained in the  $O(1)$  model can be derived writing the Hamiltonians in terms of generators of the Temperley–Lieb algebra at the semi-group point [6], acting on the link pattern (or equivalently the restricted solid on solid path) basis. Each Hamiltonian gives the Markovian time evolution of a fluctuating interface. The class of stochastic models of this type are called Raise and Peel (RP) models (see [7] for physical properties and [8] for several examples). Besides their interesting properties, these models belong to a new universality class for non-equilibrium phenomena, since the finite-size scaling spectra of these models are given by conformal field theory.

In a very recent new development, Di Francesco [9] proposed a refined RS conjecture. If one considers  $A_n(j)$  the known number [10] of  $n \times n$  ASMs with a 1 on top of the  $j$ th column one can define a generating function  $\psi_n(t)$ . According to Di Francesco’s conjecture this function coincides with the normalization of the ground state wavefunction of the  $t$  dependent monodromy matrix. As for the other cases when RS-type conjectures were made, not only the normalization, but also each component of the  $t$  dependent ground state gives information on properties of ASMs.

In this paper we will present two new refined RS-type conjectures. In section 2 we present the first conjecture. We define a Hamiltonian in terms of the  $L$  generators of the boundary Temperley–Lieb algebra. This Hamiltonian acts in the space of  $2^L$  link patterns and depends on a free parameter  $a$ . If this parameter is non-negative, the Hamiltonian gives the time evolution of a fluctuating interface. The parameter  $a$  controls the boundary rates. We have studied the ground state wavefunction, which gives the probability distribution function of the stationary state of the stochastic process, as a function of  $a$ .

For  $a = 1$ , it was observed in [11], à la Razumov and Stroganov, that the ground state wavefunction  $\psi_0(1)$  for a system of size  $L$  is related to vertically and horizontally symmetric alternating-sign matrices (VHSASMs) of size  $2L + 3$ . In the present paper we study  $\psi_0(a)$  and make a refined RS conjecture. As an application, we give the average number of  $\pm 1$ s on the  $(L + 1)$ st row of a VHSASM matrix of size  $2L + 3$ . Other results can also be used to study the properties of stationary states of the stochastic process.

Although it is not the main subject of this paper, we also comment on the connection of the Hamiltonian with the XXZ quantum chain with diagonal boundary conditions. We furthermore briefly discuss the properties of the spectrum of the Hamiltonian in the continuum limit.

The second conjecture is presented in section 3. We add to the Hamiltonian considered in section 2 a second boundary operator with a coefficient  $b$ . The new Hamiltonian depends therefore on two parameters. The space of link patterns in which the new Hamiltonian acts is different from the one in which the Hamiltonian with only one generator acts. Remarkably, the dimensions of the subspaces in which the two ground states have components are the same.

We conjecture that the ground state  $\psi_0(a, b)$  of the Hamiltonian with  $L - 1$  bulk and 2 boundary generators is related to weighted FPL diagrams on a grid of dimension  $(L + 1) \times L$  with special boundary conditions. A given configuration gets a factor  $a$  for each vertical segment at the boundary and a factor  $b$  for each horizontal segment at the boundary.

## 2. The refined one-boundary RS conjecture

The Hamiltonian that we wish to study is given by

$$H = a(1 - f_-) + \sum_{j=1}^{L-1} (1 - e_j), \quad (1)$$



**Figure 1.** An  $8 \times 2$  strip with link pattern  $)((()|()$ .

where the generators  $f_-$  and  $e_i$  satisfy the one-boundary Temperley–Lieb algebra, otherwise known as the blob algebra [12],

$$\begin{aligned}
 e_i^2 &= e_i, \\
 e_i e_{i\pm 1} e_i &= e_i, \\
 e_i e_j &= e_j e_i \quad \text{for } |i - j| \geq 2, \\
 f_-^2 &= f_-, \\
 e_1 f_- e_1 &= e_1.
 \end{aligned}
 \tag{2}$$

The Hamiltonian given by (1) describes the time evolution of a stochastic process of a fluctuating interface with a boundary; see [8] for the case  $a = 1$ . Here we are interested in the stationary state only.


The Temperley–Lieb algebra has a well known loop representation in the space of link patterns [13, 14]. Let us briefly recall this representation. Consider an  $L \times 2$  strip whose  $L$  bottom sites may be connected to each other or to the site directly above it by non-crossing arcs. In the case of the one-boundary Temperley–Lieb algebra, sites can also be connected to the left boundary of the strip. See figure 1 for an example of an  $8 \times 2$  strip. We call a particular way in which loops are connected a *link pattern* or *connectivity*, and denote the linear span of link patterns on  $L$  sites by  $LP_L$ .

On a link pattern  $\pi \in LP_L$  the generator  $e_i$  acts in the following way: if site  $i$  is connected to  $k$  and site  $i + 1$  to  $l$ ,  $e_i$  connects  $i$  with  $i + 1$  and  $k$  with  $l$ . If  $i$  is connected to  $k$  and  $i + 1$  to the top row (or left boundary) of the strip, then  $k$  gets connected to the top row (or left boundary) and  $i$  to  $i + 1$ ; similarly, if  $i$  is connected to the top row (or left boundary) and  $i + 1$  is connected to  $k$ . Lastly, if both  $i$  and  $i + 1$  are connected to the top row (or left boundary), they get connected to each other. If site 1 is connected to  $i$ , the generator  $f_-$  acts by connecting both site 1 and site  $i$  to the left boundary of the strip. If site 1 is connected to the top row it gets connected to the left boundary, while if site 1 is connected to the left boundary,  $f_-$  acts as the identity.

Link patterns can be described using a parentheses notation: if site  $i$  is connected to site  $j$  we put an opening parenthesis ‘(’ at  $i$  and a closing parenthesis ‘)’ at  $j$ . If site  $i$  is connected to the left boundary we put a closing parenthesis at  $i$ . Sites that are connected to the top of the strip are denoted by vertical bars. The link patterns for  $L = 2$  are thus given by

$$||, \quad ), \quad )), \quad (),
 \tag{3}$$

which respectively mean that (i) the two sites are connected to the top of the strip, (ii) the first is connected to the left boundary while the second is connected to the top, (iii) both are connected to the left boundary and (iv) the two sites are connected to each other.



**Figure 2.** A vertically and horizontally symmetric FPL diagram on a grid of size 9 and the corresponding alternating-sign matrix.

The dimension of the space of link patterns for the one-boundary Temperley–Lieb algebra on  $L$  sites is

$$\dim \text{LP}_L = 2^L. \quad (4)$$

Due to the semi-group structure of the algebra (2), the Hamiltonian (1) has a positive spectrum and a unique ground state energy  $E_0 = 0$  in  $\text{LP}_L$ . We will be interested in the corresponding eigenvector  $\psi_0$  as a function of the parameter  $a$ ,

$$H\psi_0(a) = 0. \quad (5)$$

This eigenvector lies in the subspace  $\text{LP}_L^0$  spanned by the link patterns without vertical bars. Its dimension is equal to

$$\dim \text{LP}_L^0 = \binom{L}{\lfloor L/2 \rfloor}. \quad (6)$$

### 2.1. Fully packed loops


In [8, 11] the Hamiltonian (1) was studied for  $a = 1$  and it was observed that  $\psi_0(1)$  for size  $L$  is related to vertically and horizontally symmetric alternating-sign matrices (VHSASMs). In particular it was conjectured that the sum of the components of  $\psi_0(1)$  for size  $L$  is equal to  $A_{\text{VH}}(2L + 3)$ , the total number of VHSASMs of size  $2L + 3$ . This number is known [15] and given by

$$A_{\text{VH}}(4n \pm 1) = A_{\text{V}}(2n \pm 1)N_8(2n), \quad (7)$$


$$A_{\text{V}}(2n + 1) = \prod_{k=0}^{n-1} (3k + 2) \frac{(6k + 3)!(2k + 1)!}{(4k + 2)!(4k + 3)!} = 1, 3, 26, 646, \dots, \quad (8)$$

$$N_8(2n) = \prod_{k=0}^{n-1} (3k + 1) \frac{(6k)!(2k)!}{(4k)!(4k + 1)!} = 1, 2, 11, 170, \dots, \quad (9)$$

where  $A_{\text{V}}(2n + 1)$  is the number of vertically symmetric  $(2n + 1) \times (2n + 1)$  alternating-sign matrices, and  $N_8(2n)$  the number of cyclically symmetric transpose complement plane partitions in a box of size  $2n \times 2n \times 2n$ . By a well known bijection [16],  $A_{\text{VH}}(2L + 3)$  is also equal to the total number of vertically and horizontally symmetric fully packed



**Figure 3.** Reduction of a  $2L + 3$  grid with vertical and horizontal symmetry to an  $L \times L$  patch.




**Figure 4.** FPL diagrams for  $L = 3$ . Diagrams with the same link pattern are grouped together. The last FPL diagram corresponds to the VHSASM given in figure 2.

loop (FPL) diagrams on grids of size  $2L + 3$ . An example of a vertically and horizontally symmetric FPL diagram on a grid of size 9 is given in figure 2. For later purposes we note that the non-zero entries in the ASM correspond to straight loop segments of length 2.

The vertical and horizontal symmetry constraint enforces certain edges to contain loop segments, and also that the four quadrants of the grid are mirror images of each other. We therefore only have to consider FPL diagrams on an  $L \times L$  patch; see figure 3. The six FPL patterns that can be drawn on the  $3 \times 3$  patch are depicted in figure 4. For each FPL diagram one can define a link pattern describing the way the external edges are connected to each other by the loop segments. We number from 1 to  $L$  the external edges that contain loop segments on the left and bottom of the  $L \times L$  patch, as in the top left diagram of figure 4. If edge  $i$  is connected to edge  $j$  we put an opening parenthesis ‘(’ at  $i$  and a closing parenthesis ‘)’ at  $j$ . If edge  $i$  is connected to one of the external edges on the top we put a closing parenthesis at  $i$ . The link patterns in figure 4 are thus  $)))$ ,  $(())$  and  $)()$  and diagrams with the same link pattern are grouped together.

Following Razumov and Stroganov [4] it was further conjectured that not only is the sum of components of  $\psi_0(1)$  related to a counting problem, but so also is each coefficient of  $\psi_0(1)$ . Writing

$$\psi_0(1) = \sum_{\pi \in \text{LP}_L^0} c_L(\pi) |\pi\rangle, \tag{10}$$



**Figure 5.** Weighted FPL diagrams for  $L = 3$ . A diagram with  $2j + 1$  vertical line segments in the top row is assigned a weight  $a^j$ . Diagrams with the same link pattern are grouped together.

the conjecture states that  $c_L(\pi)$  is equal to the number of FPL diagrams on the  $L \times L$  patch with link pattern  $\pi$ . For example for  $L = 3$  we obtain

$$\psi_0(1) = (1, 2, 3) \quad (11)$$

on the basis  $\{))) , ()), )())\}$ , indeed corresponding to the enumeration of FPL diagrams with the same link pattern; see figure 4.

## 2.2. A refined conjecture

For general values of the parameter  $a$  we find that the conjecture above is refined in the following way. If we write

$$\psi_0(a) = \sum_{\pi \in \text{LP}_L^0} c_L(\pi, a) |\pi\rangle, \quad (12)$$

the coefficient  $c_L(\pi, a)$  will be a polynomial in  $a$ . We have observed from exact calculations for  $L \leq 6$  that the coefficient of  $a^j$  in  $c_{2n}(\pi, a)$  (resp.  $c_{2n+1}(\pi, a)$ ) enumerates FPL diagrams on the  $L \times L$  patch with connectivity  $\pi$  and having  $2j$  (resp.  $2j + 1$ ) vertical line segments of length 2 measured from the top. We will illustrate this with two examples.

For  $L = 3$  we find

$$\psi_0(a) = (a, 2, 2 + a), \quad (13)$$

on the basis  $\{))) , ()), )())\}$ . Assign to each FPL diagram on the  $3 \times 3$  patch a weight  $a^j$  if it has  $2j + 1$  vertical line segments on the first row, as in figure 5. The coefficient  $c_3(\pi, a)$  of  $\psi_0(a)$  is then given by the sum of weighted FPL diagrams having link pattern  $\pi$ .

For the second example we look at the even system  $L = 4$  and obtain

$$\psi_0(a) = (a^2, 3a(2 + a), 2a(3 + a), 3a, 3(2 + a), 3), \quad (14)$$

on the basis  $\{)))) , ))(), )()), ()), (()), (())\}$ . These components are indeed recovered by the enumeration of weighted FPL diagrams, if we assign to each FPL diagram a weight  $a^j$  if it has  $2j$  vertical line segments in the top row, as in figure 6. The refined conjecture formulated above is similar to but differs from Di Francesco's recent results [9].



**Figure 6.** Weighted FPL diagrams for  $L = 4$ . A diagram with  $2j$  vertical line segments in the top row is assigned a weight  $a^j$ . Diagrams with the same link pattern are grouped together.

### 2.3. An application

The conjectures above can be used to derive properties of FPL diagrams or ASMs. For example, the normalization of the one-boundary ground state

$$Z^{(1)}(a) = \sum_{\pi \in \text{LP}_L^0} c_L(\pi, a), \tag{15}$$

is conjectured to be equal to the partition function of  $(2L + 3) \times (2L + 3)$  VHSASMs in which each  $-1$  on the row below the horizontal symmetry axis is given a weight  $a$ . The

average density  $\rho_L(a)$  of  $-1$ s on this row is thus given by

$$\rho(a) = \frac{1}{L} \frac{d \log Z^{(1)}(a)}{da}. \quad (16)$$

While we have not found a general expression for  $Z^{(1)}(a)$ , we have found numerically that

$$\rho_L(1) = \begin{cases} \frac{3L+8}{8(2L+3)} & \text{for even } L, \\ \frac{3(L^2-1)}{8L(2L+3)} & \text{for odd } L. \end{cases} \quad (17)$$

In the thermodynamic limit  $\rho_\infty(1) = 3/8$ , whereas one would expect a value of  $1/3$  in a completely random region.

#### 2.4. The relation to the XXZ chain with diagonal boundary conditions

We would like to comment on the spectrum of the Hamiltonian given in (1). It is convenient to parametrize  $a$  defined in (1) as follows:

$$a = \frac{3}{1 + 2 \cos \delta} > 0. \quad (18)$$

The spectrum of  $H$  acting in the link pattern space coincides with that of the Hamiltonian of the XXZ spin  $1/2$  quantum chain with diagonal boundary conditions and asymmetry parameter  $\Delta = -1/2$  [17]:

$$H_D = -\frac{1}{2} \left\{ \sum_{j=1}^{L-1} (\sigma_j^x \sigma_{j+1}^x + \sigma_j^y \sigma_{j+1}^y - \frac{1}{2} \sigma_j^z \sigma_{j+1}^z) + \frac{\sqrt{3}}{2} \left( \tan\left(\frac{\pi}{6} + \frac{\delta}{2}\right) (\sigma_1^z - 1) + \tan\left(\frac{\pi}{6} - \frac{\delta}{2}\right) (\sigma_L^z - 1) \right) \right\} + \frac{3}{4}(L-1), \quad (19)$$

where  $\sigma^x$ ,  $\sigma^y$  and  $\sigma^z$  are Pauli matrices. For  $a \geq 1$  ( $0 \leq \delta < 2\pi/3$ ) the Hamiltonian  $H_D$  is Hermitian. Notice that although in (1) the parameter  $a$  appears in one boundary generator only, it appears in both boundary terms in the expression for  $H_D$ . Moreover, there exists a similarity transformation which relates  $H$  to  $H_D$ . This last observation is not trivial since Jordan cell structures often appear in stochastic processes and two Hamiltonians might have the same spectrum but might not be related by a similarity transformation [17].

Since  $H$  describes a stochastic process it has a positive spectrum with a unique ground state of zero energy. The same properties are therefore inherited by  $H_D$ . In the special case  $a = 1$  ( $\delta = 0$ ), using different methods, in [18] it was shown that  $H_D$  has a positive spectrum and a zero ground state energy for all system sizes.

We will now briefly comment on the spectrum of  $H_D$  (or  $H$ ) in the continuum limit. It is independent of  $a$  [19], it is the same for  $L$  even and odd and it is given by the Gauss model. A compact way to describe the spectrum is to use [20] the sum of the characters of the  $[1/24]$  and  $[3/8]$  representations of the  $c = 1$  Ramond representations of  $N = 2$

superconformal field theory [21]. For  $a = 0$  the Hamiltonian  $H_D$  is  $U_q(sl(2))$ -invariant and the finite-size scaling spectrum is different [22].

### 3. The two-boundary case

We can extend the Hamiltonian (1) to also include a boundary term at site  $L$ ,

$$H = a(1 - f_-) + b(1 - f_+) + \sum_{j=1}^{L-1} (1 - e_j). \quad (20)$$

As before, the generators  $f_-$  and  $e_i$  satisfy the relations of the one-boundary Temperley–Lieb algebra, but with the additional relations

$$\begin{aligned} f_+^2 &= f_+, \\ e_{L-1} f_+ e_{L-1} &= e_{L-1}, \\ IJI &= I, \quad JIJ = J, \end{aligned} \quad (21)$$

where

$$I = \prod_{i=0}^{L/2-1} e_{2i+1}, \quad J = f_- \prod_{i=1}^{L/2-1} e_{2i} f_+, \quad \text{for even } L, \quad (22)$$

$$I = f_- \prod_{i=1}^{(L-1)/2} e_{2i}, \quad J = \prod_{i=1}^{(L-1)/2} e_{2i-1} f_+, \quad \text{for odd } L. \quad (23)$$

In the link pattern representation—see figure 1—loops are now also allowed to be connected to the right boundary of the strip. Connections to the right boundary are facilitated by  $f_+$  in much the same way as they are for the left boundary by  $f_-$ .

We wish to study the ground state of the Hamiltonian (20), which lies in the subspace of link patterns for which all sites are connected. One may furthermore choose not to make a distinction between the left and right boundaries, but to identify them, and this is what we will do in the following. As before we denote link patterns by sequences of parentheses and vertical bars. Parentheses are used for sites that are connected to each other while a vertical bar will now stand for sites connected to the boundary. For example, the ground state sector of the Hamiltonian (20) with identified boundaries for  $L = 4$  is given by the linear span of the six link patterns


$$|||, \quad ()|, \quad |(), \quad ||(), \quad ()(), \quad (()). \quad (24)$$

We will denote the linear span of such link patterns by  $\text{LP}_L^*$ . Its dimension is given by

$$\dim \text{LP}_L^* = \binom{L}{\lfloor L/2 \rfloor}, \quad (25)$$

which is the same as  $\dim \text{LP}_L^0$ .

We will now formulate a doubly refined conjecture based on observations for  $L \leq 6$ . For our purposes we first need to formulate a new observation: for  $a = b = 1$  and *odd*  $L$  the ground state is obtained by counting FPL configurations with appropriate link patterns



**Figure 7.** A  $6 \times 5$  patch corresponding to the ground state of the Hamiltonian (20) for  $L = 5$  and identified boundaries.

on an  $(L + 1) \times L$  patch with boundary conditions as in figure 7.<sup>3</sup> We can now formulate a doubly refined conjecture for odd systems. Writing

$$\psi_0(a, b) = \sum_{\pi \in \text{LP}_L^*} c_L(\pi, a, b) |\pi\rangle, \quad (26)$$

the coefficient  $c_L(\pi, a, b)$  is a polynomial in  $a$  and  $b$ . From exact calculations for small values of  $L = 2n + 1$  we observed that the coefficient of  $a^j b^k$  in  $c_{2n+1}(\pi, a, b)$  enumerates FPL diagrams on the  $(L + 1) \times L$  patch with connectivity  $\pi$  and having  $2j + 1$  vertical line segments in the top row and  $2k$  horizontal line segments in the last column. For  $L = 3$  we find for example

$$\psi_0(a, b) = (a + b + ab, 2 + a, 2 + b), \quad (27)$$

on the basis  $\{|||, |(), ()|\}$ , indeed corresponding to the weighted enumeration of FPL diagrams; see figure 8. In addition we note that the normalization for odd system sizes,

$$Z_{2n+1}^{(2)}(a, b) = \sum_{\pi \in \text{LP}_{2n+1}^*} c_{2n+1}(\pi, a, b), \quad (28)$$

factorizes completely,

$$Z_{2n+1}^{(2)}(a, b) = \tilde{Z}_{2n+1}^{(1)}(a) \tilde{Z}_{2n+1}^{(1)}(b). \quad (29)$$


For  $a = b = 1$  and *even*  $L$  it was conjectured in [11] that the coefficients of the ground state enumerate FPL configurations with appropriate link patterns on an  $(L + 1) \times L$  patch with boundary conditions as in figure 9. The doubly refined conjecture for even systems is formulated as follows. Write the ground state as

$$\psi_0(a, b) = \sum_{\pi \in \text{LP}_L^*} c_L(\pi, a, b) |\pi\rangle; \quad (30)$$


then the coefficient  $c_L(\pi, a, b)$  is a polynomial in  $a$  and  $b$ . We have observed from exact calculations for small values of  $L = 2n$  that the coefficient of  $a^j b^k$  in  $c_{2n}(\pi, a, b)$  enumerates FPL diagrams on the  $(L + 1) \times L$  patch with connectivity  $\pi$  and having  $2j$  or  $2j + 1$  vertical line segments in the top row and  $2k$  horizontal line segments in the last column.

We have listed the first few eigenvectors in table A.1 in the appendix.

<sup>3</sup> In [11] it was observed that the ground state is also obtained by counting FPL configurations with appropriate link patterns on an  $L \times (L + 1)/2$  patch. Compared to this observation, ours gives rise to an overall factor (equal to  $A_V(L + 2)$ ) in each of the components. The relation between the two observations is not clear.



**Figure 8.** Weighted FPL diagrams for  $L = 3$ . A diagram with  $2j + 1$  vertical line segments in the top row and  $2k$  horizontal line segments in the rightmost column is assigned a weight  $a^j b^k$ . Diagrams with the same link pattern are grouped together.



**Figure 9.** A  $7 \times 6$  patch corresponding to the ground state of the Hamiltonian (20) for  $L = 6$  and identified boundaries.

#### 4. Conclusion

This paper contains two conjectures concerning the relation between parameter dependent ground states of Hamiltonians describing stochastic processes and combinatorial properties of FPLs on grids with special boundary conditions.

We hope that the existence of more conjectures of this kind will help to produce, finally, a proof of the many known parameter-fixed conjectures which already appear in the literature (see the references mentioned in the introduction).

In both conjectures the parameters are related to boundary operators and the bookkeeping of the FPLs is also done by looking at how the loops touch the boundaries. This is in contrast to the refined conjecture of Di Francesco [9] (which was a source of inspiration for the present paper) in which the parameter changes the bulk interaction but the bookkeeping of the FPLs still counts the way the loops touch the boundary. The second difference between the two approaches can be understood in the following way.

If, in analogy with statistical physics, one interprets the parameter dependent ground state as a partition function depending on fugacities—see e.g. [23] for a justification of this interpretation—our two conjectures give information on the number of ‘defects’ on the boundary (for example the number of non-zero entries in the VHSASM matrices on a row). The refined conjecture of [9] gives the space distribution of one ‘defect’ (the position of 1 in the first row of ASMs).

## Acknowledgments

We would like to thank A Nichols for discussions and G Kuperberg for useful correspondence. This work was done within the European Commission network HPRN-CT-2002-00325. Financial support from the Deutsche Forschungsgemeinschaft and the Australian Research Council is gratefully acknowledged.

## Appendix. The eigenvector $\psi_0$

In this appendix we list the eigenvectors for the two-boundary case and  $L \leq 4$ . The eigenvectors for one boundary are obtained by setting  $b = 0$ .

**Table A.1.** The first few eigenvectors for the two-boundary case.

$L$	Basis	$\psi_0$
1	{ }	{1}
2	{  , ()}	{ $a + b, 1$ }
3	{   ,  (), ()}	{ $a + b + ab, 2 + a, 2 + b$ }
4	{    ,   (),  (), () , ()(), ()}	{ $a^2 + 4ab + 2a^2b + b^2 + 2ab^2, 3(2a + a^2 + b + ab),$ $6a + 2a^2 + 6b + 5ab + a^2b + 2b^2 + ab^2, 3(a + 2b + ab + b^2),$ $3(2 + a + b), 3$ }

## References

- [1] Stroganov Yu G, *The importance of being odd*, 2001 *J. Phys. A: Math. Gen.* **34** L179  
Razumov A V and Stroganov Yu G, *Spin chains and combinatorics*, 2001 *J. Phys. A: Math. Gen.* **34** 3185
- [2] Bressoud D, 1999 *Proofs and Confirmations. The Story of the Alternating Sign Matrix Conjecture* (Cambridge: Cambridge University Press)
- [3] Batchelor M T, de Gier J and Nienhuis B, *The quantum symmetric XXZ chain at  $\Delta = -1/2$ , alternating sign matrices and plane partitions*, 2001 *J. Phys. A: Math. Gen.* **34** L265
- [4] Razumov A V and Stroganov Yu G, *Combinatorial nature of ground state vector of  $O(1)$  loop model*, 2001 *Theor. Math. Phys.* **138** 333 [[math.CO/0104216](#)]
- [5] Razumov A V and Stroganov Yu G, *Spin chains and combinatorics: twisted boundary conditions*, 2001 *J. Phys. A: Math. Gen.* **34** 5335 [[cond-mat/0102247](#)]  
Razumov A V and Stroganov Yu G,  *$O(1)$  loop model with different boundary conditions and symmetry classes of alternating-sign matrices*, 2001 Preprint [cond-mat/0108103](#)
- [6] Pearce P A, Rittenberg V, de Gier J and Nienhuis B, *Temperley–Lieb stochastic processes*, 2002 *J. Phys. A: Math. Gen.* **35** L661 [[math-ph/0209017](#)]
- [7] de Gier J, Nienhuis B, Pearce P A and Rittenberg V, *Stochastic processes and conformal invariance*, 2003 *Phys. Rev. E* **67** 016101  
de Gier J, Nienhuis B, Pearce P A and Rittenberg V, *The Raise and Peel model of a fluctuating interface*, 2004 *J. Stat. Phys.* **114** 1

- [8] Pyatov P, *Raise and Peel models of fluctuating interfaces and combinatorics of Pascal’s hexagon*, 2004 *J. Stat. Mech.: Theor. Exp.* P09003 [[math-ph/0406025](#)]
- [9] Di Francesco P, *A refined Razumov–Stroganov conjecture*, 2004 *J. Stat. Mech.: Theor. Exp.* P08009 [[cond-mat/0407477](#)]
- [10] Zeilberger D, *Proof of the refined alternating sign matrix conjecture*, 1996 *New York J. Math.* **2** 59
- [11] Mitra S, Nienhuis B, de Gier J and Batchelor M T, *Exact expressions for correlations in the ground state of the dense  $O(1)$  loop model*, 2004 *J. Stat. Mech.: Theor. Exp.* to be published [[cond-mat/0401245](#)]
- [12] Martin P P and Saleur H, *The blob algebra and the periodic Temperley–Lieb algebra*, 1994 *Lett. Math. Phys.* **30** 189
- [13] Temperley H N V and Lieb E H, *Relations between the ‘percolation’ and ‘colouring’ problem and other graph-theoretical problems associated with regular planar lattices: some exact results for the ‘percolation’ problem*, 1971 *Proc. R. Soc. A* **322** 251
- [14] Martin P P, 1991 *Potts Models and Related Problems in Statistical Mechanics* (Singapore: World Scientific)
- [15] Okada S, *Enumeration of symmetry classes of alternating sign matrices and characters of classical groups*, 2004 Preprint [math.CO/0408234](#) (<http://www.maths.anu.edu.au/events/CIM/Slides/Okada.pdf>)  
Robbins D P, *Symmetry classes of alternating sign matrices*, 2000 Preprint [math.CO/0008045](#)  
Kuperberg G, *Symmetry classes of alternating-sign matrices under one roof*, 2002 *Ann. Math.* **156** 835 [[math.CO/0008184](#)]
- [16] Batchelor M T, Blöte H W J, Nienhuis B and Yung C M, *Critical behaviour of the fully packed loop model on the square lattice*, 1996 *J. Phys. A: Math. Gen.* **29** L399  
Propp J, *The many faces of alternating-sign matrices*, 2001 *Discrete Mathematics and Theoretical Computer Science Proceedings AA* p 43 (<http://dmtcs.loria.fr/proceedings/dmAAind.html>)
- [17] Nichols A, private communication
- [18] Fendley P, Schoutens K and Nienhuis B, *Lattice fermion models with supersymmetry*, 2002 *J. Phys. A: Math. Gen.* **36** 12399  
Yang X and Fendley P, *Non-local space-time supersymmetry on the lattice*, 2004 Preprint [cond-mat/0404682](#)  
Beccaria M and De Angelis G F, *Exact ground state and finite size scaling in a supersymmetric lattice model*, 2004 Preprint [cond-mat/0407752](#)
- [19] Alcaraz F C, Barber M N and Batchelor M T, *Conformal invariance and the spectrum of the XXZ chain*, 1987 *Phys. Rev. Lett.* **58** 771  
Alcaraz F C, Baake M, Grimm U and Rittenberg V, *The modified XXZ Heisenberg chain, conformal invariance and the surface exponents of  $c < 1$  systems*, 1989 *J. Phys. A: Math. Gen.* **22** L5  
Grimm U and Rittenberg V, *The modified XXZ Heisenberg chain, conformal invariance, surface exponents of  $c < 1$  systems, and hidden symmetries of the finite chains*, 1990 *Int. J. Mod. Phys. B* **4** 669
- [20] de Gier J, Nichols A, Pyatov P and Rittenberg V, 2004 in preparation
- [21] Ravanini F and Yang S-K,  *$C$ -disorder fields and  $\Gamma(2)$ -invariant partition functions in parafermionic conformal field theories*, 1988 *Nucl. Phys. B* **295** 262  
Eguchi T and Yang S-K,  *$N = 2$  superconformal models as topological field theories*, 1990 *Mod. Phys. Lett.* **5** 1693  
Saleur H and Warner N P, *Lattice models and  $N = 2$  supersymmetry*, 1993 Preprint [hep-th/9311138](#)
- [22] Bauer M and Saleur H, *On some relations between local height probabilities and conformal invariance*, 1989 *Nucl. Phys. B* **320** 591
- [23] Brak R, de Gier J and Rittenberg V, *Nonequilibrium stationary states and equilibrium models with long range interactions*, 2004 *J. Phys. A: Math. Gen.* **37** 4303