

ON INTEGRAL REPRESENTATIONS OF THE DRAZIN INVERSE IN BANACH ALGEBRAS

by N. CASTRO GONZÁLEZ, J. J. KOLIHA and YIMIN WEI*

The purpose of the present paper is to derive an integral representation of the Drazin inverse of an element of a Banach algebra in a more general situation than previously obtained by the second author, and to give an application to the Moore–Penrose inverse in a C^* -algebra.

2000 *Mathematics subject classification*: 46H05, 46L05

1 Introduction

Let \mathcal{A} be a complex unital Banach algebra with unit e . In [4], a generalized *Drazin inverse* of an element $a \in \mathcal{A}$ was defined as $b \in \mathcal{A}$ such that

$$ab = ba, \quad b^2a = b, \quad a^2b = a + u, \quad (1.1)$$

where $u \in \mathcal{A}$ is quasinilpotent, that is, $\lim_{n \rightarrow \infty} \|u^n\|^{1/n} = 0$ [4, Definition 4.1]. (See also [3].) This definition subsumes (for Banach algebras) the pseudo-inverse defined originally for elements of semigroups and rings [2], which arises when u is nilpotent. The Drazin inverse b of a is unique when it exists, and is denoted by a^D . The *Drazin index* $i(a)$ of a is defined to be 0 if a is invertible, k if the element u in (1.1) is nilpotent of order k , and ∞ otherwise.

According to [4], an element $a \in \mathcal{A}$ is Drazin invertible if and only if 0 is not an accumulation point of $\sigma(a)$. This occurs if and only if there exists an idempotent $p \in \mathcal{A}$ such that

$$ap = pa \text{ is quasinilpotent, } \quad a + p \text{ is invertible} \quad (1.2)$$

[4, Theorem 4.2]; p is the *spectral idempotent* of a denoted by a^π . We have

$$a^D = (a + a^\pi)^{-1}(e - a^\pi) \quad \text{and} \quad a^\pi = e - a^D a. \quad (1.3)$$

*Project 19901006 supported by the National Natural Science Foundation of China and China Scholarship Council.

We also need the core–quasinilpotent decomposition of a Drazin invertible element $a \in \mathcal{A}$ introduced in [4] in the form $a = x + y$ where $xy = yx = 0$, x is of the Drazin index not exceeding 1, and y is quasinilpotent; x is called the *core* of a . Explicitly, $x = a(e - a^\pi)$. The importance of the core of a is reflected in the equations

$$i(x) \leq 1, \quad \sigma(x) = \sigma(a), \quad x^D = a^D. \quad (1.4)$$

Various representation of the Drazin inverse, mostly for matrices, appear in the literature. (See, for instance, [8, 10, 11].)

In [4], an integral representation was given for an element $a \in \mathcal{A}$ for which $\exp(ta)$ converges as $t \rightarrow \infty$. This representation turned out to be a useful tool in theory of singular differential equations, where it was applied to derive conditions for the asymptotic convergence of solutions both in the setting of matrices [6] and semigroups of operators [1, 7].

The purpose of the present paper is to derive an integral representation of the Drazin inverse in a more general situation than in [4] and give an application to the Moore–Penrose inverse in a C^* -algebra.

2 The integral representation

We say that $a \in \mathcal{A}$ is *semistable* if a is Drazin invertible with $\text{ind}(a) \leq 1$ and the nonzero spectrum of a lies in the open left half of the complex plane. The following result is [4, Theorem 6.3].

Proposition 2.1. *Let $a \in \mathcal{A}$ be semistable with the spectral idempotent a^π . Then*

$$a^D = - \int_0^\infty \exp(ta)(e - a^\pi) dt. \quad (2.1)$$

In our first main result we show that the integral representation remains true for a with an arbitrary Drazin index.

Theorem 2.2. *Let $a \in \mathcal{A}$ be a Drazin invertible element with a finite or infinite Drazin index such that the nonzero spectrum of a lies in the open left half of the complex plane. Then equation (2.1) holds.*

Proof. The hypothesis of the Drazin invertibility of a implies that 0 is a resolvent point or an isolated spectral point of a . Let $p = a^\pi$ and let $x = a(e - p)$ be the

core of a . In view of (1.4), x is semistable, and $x^D = -\int_0^\infty \exp(tx)(e-p) dt$ by Proposition 2.1. Further,

$$\begin{aligned} \exp(tx)(e-p) &= \exp(ta(e-p))(e-p) = (p + \exp(ta)(e-p))(e-p) \\ &= \exp(ta)(e-p), \end{aligned}$$

and

$$a^D = x^D = -\int_0^\infty \exp(tx)(e-p) dt = -\int_0^\infty \exp(ta)(e-p) dt. \quad \square$$

The following representation is valid for elements of finite Drazin index.

Theorem 2.3. *Let $a \in \mathcal{A}$ be a Drazin invertible element with a finite Drazin index $k \geq 1$ such that for some $n \geq 1$ the nonzero spectrum of a^n lies in the open left half of the complex plane. Then, for any $m \geq k$,*

$$-\int_0^\infty \exp(ta^n)a^m dt = (a^D)^n a^m = \begin{cases} (a^D)^{n-m} & \text{if } m < n, \\ e - a^\pi & \text{if } m = n, \\ x^{m-n} & \text{if } m > n. \end{cases} \quad (2.2)$$

Proof. Let $a \in \mathcal{A}$ be a Drazin invertible element with $p = a^\pi$. Then a^n is also Drazin invertible, and $(a^n)^D = (a^D)^n$ [4, Theorem 5.4]. In view of (1.3), the spectral idempotent of a^n is also equal to p :

$$e - (a^n)^D a^n = e - (a^D a)^n = e - a^D a = p.$$

Applying Theorem 2.2 to a^n in place of a and using equation $pa^m = 0$, we get

$$\int_0^\infty \exp(ta^n)a^m dt = \int_0^\infty \exp(ta^n)(e-p)a^m dt = -(a^n)^D a^m. \quad (2.3)$$

By (1.3) again,

$$(a^D)^n a^m = (a+p)^{-n}(e-p)(a+p)^m = (a+p)^{m-n}(e-p),$$

from which (2.2) follows when we observe that $x^r = a^r(e-p)$ for any $r > 0$. \square

Specializing the preceding theorem, we get a new integral representation for the Drazin inverse.

Theorem 2.4. *Let $a \in \mathcal{A}$ be a Drazin invertible element of a finite Drazin index $k \geq 1$ such that the nonzero spectrum of a^{m+1} lies in the open left half of the complex plane for some $m \geq k$. Then*

$$a^{\text{D}} = - \int_0^{\infty} \exp(ta^{m+1})a^m dt. \quad (2.4)$$

The condition that the nonzero spectrum of a^{m+1} lies in the open left half of the complex plane is equivalent to the condition that the nonzero spectrum of a lies in the union of $m + 1$ angular regions

$$\frac{4j+1}{2(m+1)}\pi < \theta < \frac{4j+3}{2(m+1)}\pi, \quad j = 0, 1, \dots, m.$$

(Divide the unit ‘pie’ into $2(m+1)$ equal slices starting at $\theta = \pi/(2(m+1))$ and keep every second slice starting with $\pi/(2(m+1)) < \theta < 3\pi/(2(m+1))$.)

There is also a ‘right half plane’ version of Theorem 2.4.

Corollary 2.5. *Let $a \in \mathcal{A}$ be a Drazin invertible element of a finite Drazin index $k \geq 1$ such that the nonzero spectrum of a^{m+1} lies in the open right half of the complex plane for some $m \geq k$. Then*

$$a^{\text{D}} = \int_0^{\infty} \exp(-ta^{m+1})a^m dt. \quad (2.5)$$

3 Application to Moore–Penrose inverse

Let \mathcal{A} be a unital C^* -algebra. According to [5, Theorem 2.5], $a \in \mathcal{A}$ is *Moore–Penrose invertible* if and only if a^*a (respectively aa^*) is Drazin invertible with the Drazin index not exceeding 1. We observe that

$$ap = 0 = pa, \quad (3.1)$$

where p is the (self-adjoint) spectral idempotent of a^*a (and also of aa^*):

$$\|ap\|^2 = \|(ap)^*ap\| = \|pa^*ap\| = 0, \quad \|pa\|^2 = \|pa(pa)^*\| = \|paa^*p\| = 0.$$

The *Moore–Penrose inverse* of a can be then defined by

$$a^\dagger = (a^*a)^{\text{D}}a^* = a^*(aa^*)^{\text{D}}. \quad (3.2)$$

Since the nonzero spectrum of a^*a always lies in the open right half of the complex plane and the Drazin index of a^*a does not exceed 1, Proposition 2.1 and Corollary 2.5 apply to give the following representation of the Moore–Penrose inverse.

Theorem 3.1. *Let $a \in \mathcal{A}$ be a Moore–Penrose invertible element of a C^* -algebra \mathcal{A} . Then, for each $m \geq 0$,*

$$a^\dagger = \int_0^\infty \exp(-t(a^*a)^{m+1})(a^*a)^m a^* dt = \int_0^\infty a^* \exp(-t(aa^*)^{m+1})(aa^*)^m dt. \quad (3.3)$$

Proof. Equation (3.3) with $m = 0$ is obtained when we apply Proposition 2.1 to the formula (3.2) for the Moore–Penrose inverse, taking into account that $a^*p = pa^* = 0$ in view of (3.1). We have thus obtained Showalter’s representation [9] by a different method.

The case $m > 0$ follows from Corollary 2.5. □

References

- [1] N. Castro González and J. J. Koliha, Perturbation of the Drazin inverse for closed linear operators, *Integral Equations Operator Theory* **36** (2000), 92–106.
- [2] M. P. Drazin, Pseudo-inverse in associative rings and semigroups, *Amer. Math. Monthly* **65** (1958), 506–514.
- [3] R. E. Harte, Spectral projections, *Irish Math. Soc. Newsletter* **11** (1984), 10–15.
- [4] J. J. Koliha, A generalized Drazin inverse, *Glasgow Math. J.* **38** (1996), 367–381.
- [5] J. J. Koliha, The Drazin and Moore–Penrose inverse in C^* -algebras, *Math. Proc. Roy. Irish Acad.* **99A** (1999), 17–27.
- [6] J. J. Koliha and Ivan Straškraba, Power bounded and exponentially bounded matrices, *Appl. Math.* **44** (1999), 289–308.
- [7] J. J. Koliha and T. D. Tran, The Drazin inverse for closed linear operators, *J. Operator Theory*, to appear.
- [8] U. G. Rothblum, A representation of the Drazin inverse and characterization of the index, *SIAM J. Appl. Math.* **31** (1976), 646–648.
- [9] D. Showalter, Representation and computation of the pseudoinverse, *Proc. Amer. Math. Soc.* **18** (1967), 584–586.
- [10] Yimin Wei, A characterization and representation of the Drazin inverse, *SIAM J. Matrix Anal. Appl.* **17** (1996), 744–747.
- [11] Yimin Wei and Hebing Wu, The representation and approximation of the Drazin inverse, *J. Comp. Appl. Math.* **126** (2000), 417–432.

DEPARTAMENTO DE MATEMÁTICA APLICADA
FACULTAD DE INFORMÁTICA
UNIVERSIDAD POLITÉCNICA DE MADRID
SPAIN
nieves@fi.upm.es

DEPARTMENT OF MATHEMATICS
AND STATISTICS
UNIVERSITY OF MELBOURNE
AUSTRALIA
j.koliha@ms.unimelb.edu.au

DEPARTMENT OF MATHEMATICS
FUDAN UNIVERSITY
SHANGHAI 200433
P. R. CHINA
ymwei@fudan.edu.cn