

ON THE ORIENTED CHROMATIC NUMBER OF DENSE
GRAPHS

DAVID R. WOOD

ABSTRACT. Let G be a graph with n vertices, m edges, average degree d , and maximum degree Δ . The *oriented chromatic number* of G is the maximum, taken over all orientations of G , of the minimum number of colours in a proper vertex colouring such that between every pair of colour classes all edges have the same orientation. We investigate the oriented chromatic number of graphs, such as the hypercube, for which $d \geq \log n$. We prove that every such graph has oriented chromatic number at least $\Omega(\sqrt{n})$. In the case that $d \geq (2 + \epsilon) \log n$, this lower bound is improved to $\Omega(\sqrt{m})$. Through a simple connection with harmonious colourings, we prove a general upper bound of $\mathcal{O}(\Delta\sqrt{n})$ on the oriented chromatic number. Moreover this bound is best possible for certain graphs. These lower and upper bounds are particularly close when G is $(c \log n)$ -regular for some constant $c > 2$, in which case the oriented chromatic number is between $\Omega(\sqrt{n \log n})$ and $\mathcal{O}(\sqrt{n \log n})$.

1. INTRODUCTION

Throughout this paper, G is a finite simple undirected graph with n vertices, m edges, and maximum degree Δ . No loops and no parallel edges are allowed. A *colouring* of G is a function $c : V(G) \rightarrow X$, for some set of ‘colours’ X , such that $c(v) \neq c(w)$ for each edge $vw \in E(G)$. The *chromatic number* of G , denoted by $\chi(G)$, is the minimum number of colours in a colouring of G . An *orientation* of G is a directed graph D obtained from G by giving each edge a direction. Then D is called an *oriented graph*. An *oriented colouring* of an oriented graph D is a colouring c of the underlying undirected graph of D , such that between each pair of colour classes, all edges have the same direction; that is, there are no arcs \vec{vw} and \vec{xy} in D with $c(v) = c(y)$ and $c(w) = c(x)$. The *oriented chromatic number* of D , denoted by $\vec{\chi}(D)$, is the minimum number of colours in an oriented colouring of D . The *oriented chromatic number* of an undirected graph G , denoted

Received by the editors February 22, 2007, and in revised form August 10, 2007.

2000 *Mathematics Subject Classification*. 05C15.

Key words and phrases. graph, graph colouring, oriented colouring, oriented chromatic number, hypercube, harmonious colouring, degree / diameter problem.

Research supported by a Marie Curie Fellowship of the European Commission under contract MEIF-CT-2006-023865, and by the projects MEC MTM2006-01267 and DURSI 2005SGR00692.

by $\vec{\chi}(G)$, is the maximum of $\vec{\chi}(D)$, taken over all orientations D of G . The oriented chromatic number was introduced in [9] in 1994 and is now a widely studied parameter; see [3, 4, 5, 6, 7, 9, 10, 11, 17, 19, 21, 25, 26, 27, 28, 29, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 47].

This paper is motivated by a question of André Raspaud [private communication, Prague 2004], who asked for the oriented chromatic number of the d -dimensional *hypercube* Q_d . This is the graph with vertex set $\{0, 1\}^d$, where two vertices are adjacent whenever they differ in precisely one coordinate. Q_d is d -regular and has 2^d vertices. In this paper we prove generally applicable bounds on $\vec{\chi}(G)$, which in the case of the hypercube give

$$(1.1) \quad 0.8007\dots \sqrt{2^d} \leq \vec{\chi}(Q_d) \leq 2d\sqrt{2^d - 1},$$

thus determining $\vec{\chi}(Q_d)$ to within a factor of about $(5/2)d$. No non-trivial bounds on $\vec{\chi}(Q_d)$ were previously known, as we now describe.

An undirected graph has $\chi(G) = n$ if and only if $G = K_n$ if and only if G has diameter 1. But when does $\vec{\chi}(G) = n$? This question was asked by [21], who observed that for every oriented graph D ,

$$(1.2) \quad \vec{\chi}(D) = n \text{ if and only if } D \text{ has diameter } 2.$$

Here the *diameter* of D is the least integer k such that every (unordered) pair of vertices in D are connected by a directed path of at most k edges. Klostermeyer and MacGillivray [25] call an oriented graph with diameter 2 an *oclique*. Note that small diameter (> 2) does not necessarily imply large oriented chromatic number. For example, $K_{1,1,n}$ has an orientation with diameter 3 and oriented chromatic number 3. Erdős [16] proposed studying the extremal function $f(n)$, defined to be the minimum number of arcs in an oriented graph with n vertices and diameter 2. Katona and Szemerédi [24] prove that $(n/2) \log(n/2) \leq f(n) \leq n \lceil \log n \rceil$. Füredi et al. [21] tighten both bounds to conclude that $f(n) = (1 - o(1))n \log n$. The same result is independently obtained in [2]. These results imply that there are n -vertex graphs with approximately the same number of edges as the hypercube (that is, $n \log n$), yet have oriented chromatic number n . Thus good bounds for $\vec{\chi}(Q_d)$ cannot be obtained solely in terms of the number of edges.

The example of an oriented graph with diameter 2 by Füredi et al. [21] has a vertex of degree $n-1$. Thus it is natural to consider the oriented chromatic number of graphs with bounded degree. Sopena [42] and Kostochka et al. [29] prove that the oriented chromatic number is bounded for graphs of bounded degree. The best bound is due to Kostochka et al. [29], who prove that every graph G satisfies

$$(1.3) \quad \vec{\chi}(G) \leq 2\Delta^2 2^\Delta,$$

and if G is Δ -regular with sufficiently many vertices then

$$(1.4) \quad \vec{\chi}(G) \geq 2^{\Delta/2}.$$

Thus the exponential dependence on Δ in (1.3) is unavoidable. Observe that for graphs such as the hypercube with $\Delta \geq \log n$, the upper bound in (1.3) is greater than the trivial upper bound of n .

This motivates the study of the oriented chromatic number of graphs whose average degree is at least logarithmic in the number of vertices. For any such graph we establish a lower bound of $\Omega(\sqrt{n})$ on the oriented chromatic number. If the average degree is at least $(2 + \epsilon) \log n$ then this lower bound is improved to $\Omega(\sqrt{m})$. These results are proved in Section 4. In Section 2 we use a simple connection with harmonious colourings to prove a general upper bound of $\mathcal{O}(\Delta\sqrt{n})$ on the oriented chromatic number. Moreover this bound is best possible for certain graphs, as proved in Section 3.

2. AN UPPER BOUND

In this section we prove an elementary upper bound on the oriented chromatic number. A colouring of an undirected graph G is *harmonious* if the endpoints of every pair of distinct edges receive at least three colours. That is, every bichromatic subgraph has at most one edge. The *harmonious chromatic number* $h(G)$ is the minimum number of colours in a harmonious colouring of G ; see the survey [12] and the more recent references [8, 13, 14, 15, 22, 23, 30, 46].

The following two basic lower bounds on $h(G)$ are well known. Since a vertex and its neighbours all receive distinct colours in a harmonious colouring,

$$h(G) \geq \Delta + 1.$$

Since G has at most $\binom{h(G)}{2}$ edges,

$$(2.1) \quad h(G) > \sqrt{2m}.$$

The next bound is new. Observe that a harmonious colouring of G is an oriented colouring for every orientation of G . Thus

$$(2.2) \quad \vec{\chi}(G) \leq h(G).$$

Many upper bounds on $h(G)$ are known. For example, in [32] it is proved that $h(G) \leq 2\Delta\sqrt{n-1}$. Thus (2.2) implies the following lemma, which proves the upper bound on $\vec{\chi}(Q_d)$ in (1.1).

Lemma 2.1. *For every graph G ,*

$$\vec{\chi}(G) \leq 2\Delta\sqrt{n-1}.$$

3. AN EXISTENTIAL LOWER BOUND

In this section we construct a graph whose oriented chromatic number is within a constant factor of the upper bound in Lemma 2.1. To do so we construct an n -vertex graph with diameter 2 and small maximum degree Δ . Then by (1.2), the oriented chromatic number will be n . Minimising Δ is a special case of the degree/diameter problem, which asks for large graphs with given diameter and given degree; see [33] for a survey. As far as we

are aware, this variant of the degree/diameter problem has not been studied previously. (Numerous papers consider the degree/diameter problem for directed graphs with antiparallel edges.) By Moore's bound for undirected graphs, we have $\Delta \geq \sqrt{n-1}$. We now prove that, up to a multiplicative constant, this lower bound can be attained by a construction.

Lemma 3.1. *For infinitely many n , there is an n -vertex Δ -regular graph G with $\vec{\chi}(G) = n$ and $\Delta = \sqrt{8n+1} - 3$.*

Proof. As illustrated in Figure 1, let G be the underlying undirected graph of the oriented graph D with vertex set $\{(i, j) : 1 \leq i < j \leq p\}$ and arcs

- (a) $(i, j)(i, k)$ whenever $i < j < k$,
- (b) $(i, j)(k, j)$ whenever $i < k < j$, and
- (c) $(i, j)(k, i)$ whenever $k < i < j$.




FIGURE 1. Construction of the oriented graph D .

First we compute the degree of each vertex (i, j) . Observe that (i, j) has $p-j$ outgoing type-(a) arcs, $j-i-1$ outgoing type-(b) arcs, and $i-1$ outgoing type-(c) arcs. Thus (i, j) has outdegree $(p-j) + (j-i-1) + (i-1) = p-2$. A type-(a) incoming arc at (i, j) is from a vertex (i, k) with $i < k < j$; there are $j-i-1$ such arcs. A type-(b) incoming arc at (i, j) is from a vertex (k, j) with $k < i < j$; there are $i-1$ such arcs. A type-(c) incoming arc at (i, j) is from a vertex (j, k) with $i < j < k$; there are $p-j$ such arcs. Thus (i, j) has indegree $(j-i-1) + (i-1) + (p-j) = p-2$. Hence G is Δ -regular with $\Delta = 2(p-2) = \sqrt{8n+1} - 3$.

Suppose on the contrary that D has a directed 2-cycle C . If C has a type-(a) arc $(i, j)(i, k)$, then the reverse arc $(i, k)(i, j)$ is also type-(a), implying $j < k$ and $k < j$, which is a contradiction. If C has a type-(b) arc $(i, j)(k, j)$, then the reverse arc $(k, j)(i, j)$ is also type-(b), implying $k < j$ and $j < k$, which is a contradiction. If C has a type-(c) arc $(i, j)(k, i)$, then the reverse arc is $(k, i)(i, j)$, but there are no arcs of this form. Thus D has no directed 2-cycle, and indeed D is an oriented graph.

We claim that D has diameter 2. Consider two vertices (i, j) and (k, ℓ) . Then $i < j$ and $k < \ell$, and without loss of generality, $i \leq k$. If $i = k$ and $j < \ell$, then $(i, j)(i, \ell)$ is a type-(a) arc of D . If $i = k$ and $\ell < j$, then

$(i, \ell)(i, j)$ is a type-(a) arc of D . Now assume that $i < k$. If $i < k$ and $j = \ell$, then $(i, j)(k, j)$ is a type-(b) arc of D . If $i < k$ and $j < \ell$, then $(i, j)(i, \ell)(k, \ell)$ is a type-(ab) path of D . Otherwise $i < k$ and $\ell < j$, implying $i < k < \ell < j$, in which case $(i, j)(\ell, j)(k, \ell)$ is a type-(bc) path of D . Thus D has diameter 2, implying $\vec{\chi}(D) = n$ by (1.2). \square

It follows from Lemma 3.1 that in any upper bound of the form $\vec{\chi}(G) \leq \mathcal{O}(\Delta^\alpha n^\beta)$, we must have $\alpha + 2\beta \geq 2$. In particular with $\beta = 1/2$, the graph G from Lemma 3.1 has $\vec{\chi}(G) = n > \Delta\sqrt{n/8}$. In this sense, the upper bound in Lemma 2.1 is tight up to a constant factor.

4. A UNIVERSAL LOWER BOUND

We now consider universal lower bounds on the oriented chromatic number. Kostochka et al. [29] prove the following¹ lower bound for all G , which implies (1.4). (Throughout this paper, all logarithms are base 2.)

$$(4.1) \quad \binom{\vec{\chi}(G)}{2} + n \log(\vec{\chi}(G)) \geq m.$$

We now reformulate (4.1) for reasonably dense graphs. Say G has average degree $d := 2m/n$. Let t be the solution to

$$(4.2) \quad t + \log t = d - \log n.$$

Note that $0 < t < d$ and $t \rightarrow d$ for $d \gg \log n$. (We are not interested in the case $d \ll \log n$, when t becomes small.)

Lemma 4.1. *For every graph G , where t is defined as in (4.2),*

$$\vec{\chi}(G) \geq \sqrt{nt}.$$

Proof. Suppose to the contrary that $\vec{\chi}(G) < \sqrt{nt}$. By (4.1),

$$\binom{\sqrt{nt}}{2} + n \log(\sqrt{nt}) > m.$$

Thus

$$\frac{nt}{2} + \frac{n \log(nt)}{2} > \frac{dn}{2},$$

implying $t + \log t > d - \log n$. This contradiction proves the claim. \square

Lemma 4.2. *For every graph G with average degree $d \geq \log n$,*

$$\vec{\chi}(G) \geq 0.8007\dots\sqrt{n}.$$

Proof. Lemma 4.1 implies the claim since $\sqrt{t} \geq 0.8007\dots$ whenever $d \geq \log n$. \square

¹For completeness we include the proof of Equation (4.1) in [29]. Let $k := \vec{\chi}(G)$. G has less than k^n colourings with k colours, each of which is an oriented colouring of at most $2^{\binom{k}{2}}$ orientations. Thus the number of orientations, 2^m , is less than $k^n 2^{\binom{k}{2}}$. Thus $m < n \log k + \binom{k}{2}$.

For the hypercube, $d = \log n$. Thus Lemma 4.2 implies the lower bound in (1.1). Since (4.1) is proved by a non-constructive counting argument, it would be interesting to construct an orientation D of Q_d with $\vec{\chi}(D) \in \Omega(\sqrt{2^d})$; see [1, 18, 20, 31] for results on specific orientations of the hypercube.

We now refine Lemma 4.1 for graphs that are more dense than hypercubes.

Lemma 4.3. *For every graph G with average degree $d \geq \log n + (1 + \epsilon) \log t$ for some $\epsilon > 0$, where t is defined as in (4.2),*

$$\vec{\chi}(G) \geq \sqrt{\frac{\epsilon}{1 + \epsilon} (2m - n \log n)}.$$

(For example, the assumption in Lemma 4.3 holds if $d \geq (2 + \epsilon) \log n$.)

Proof. By the assumption, $t + \log t \geq (1 + \epsilon) \log t$ and $t > \epsilon \log t$. Thus

$$(1 + \epsilon)t > \epsilon(t + \log t) = \epsilon(d - \log n)$$

and

$$(1 + \epsilon)tn > \epsilon(dn - n \log n) = \epsilon(2m - n \log n).$$

Therefore Lemma 4.1 implies that

$$\vec{\chi}(G) \geq \sqrt{tn} > \sqrt{\frac{\epsilon}{1 + \epsilon} (2m - n \log n)}.$$

□

Lemma 4.3 says that for sufficiently dense graphs (that is, graphs with super-logarithmic average degree) the lower bound of $h(G) \geq \Omega(\sqrt{m})$ in (2.1) also holds for the oriented chromatic number.

Now suppose that G is Δ -regular for some $\Delta \geq (2 + \epsilon) \log n$. Thus Lemmas 2.1 and 4.3 determine $\vec{\chi}(G)$ to within a factor of $\Theta(\sqrt{\Delta})$. In particular,

$$(4.3) \quad \sqrt{\frac{\epsilon}{2 + \epsilon} \Delta n} \leq \sqrt{\frac{\epsilon}{1 + \epsilon} (\Delta - \log n)n} \leq \vec{\chi}(G) \leq 2\Delta\sqrt{n - 1}.$$

The bounds in (4.3) are particularly close when G is $(c \log n)$ -regular for some constant $c > 2$. Then $\vec{\chi}(G)$ is between $\Omega(\sqrt{n \log n})$ and $\mathcal{O}(\sqrt{n \log n})$.

ACKNOWLEDGEMENTS

Thanks to Vida Dujmović, André Raspaud, Bruce Reed, Jéan-Sebastien Sereni, Ricardo Strausz, and Stéphan Thomassé for stimulating discussions.

REFERENCES

1. Pierre Baldi, *Neural networks, orientations of the hypercube, and algebraic threshold functions*, IEEE Trans. Inform. Theory **34** (1988), no. 3, 523–530.
2. Béla Bollobás and Alex Scott, *Separating systems and oriented graphs of diameter two*, J. Combin. Theory Ser. B **97** (2007), no. 2, 193–203.
3. Oleg V. Borodin, Dmitry Fon-Der-Flaass, Alexandr V. Kostochka, André Raspaud, and Éric Sopena, *On deeply critical oriented graphs*, J. Combin. Theory Ser. B **81** (2001), no. 1, 150–155.

4. Oleg V. Borodin and A. O. Ivanova, *Oriented 7-coloring of planar graphs with girth at least seven*, Sib. Èlektron. Mat. Izv. **2** (2005), 222–229.
5. ———, *Oriented coloring of planar graphs with girth at least four*, Sib. Èlektron. Mat. Izv. **2** (2005), 239–249.
6. Oleg V. Borodin, Alexandr V. Kostochka, Jaroslav Nešetřil, André Raspaud, and Éric Sopena, *On universal graphs for planar oriented graphs of a given girth*, Discrete Math. **188** (1998), no. 1–3, 73–85.
7. ———, *On the maximum average degree and the oriented chromatic number of a graph*, Discrete Math. **206** (1999), no. 1–3, 77–89.
8. Duncan Campbell and Keith Edwards, *A new lower bound for the harmonious chromatic number*, Australas. J. Combin. **29** (2004), 99–102.
9. Bruno Courcelle, *The monadic second order logic of graphs VI: On several representations of graphs by relational structures*, Discrete Appl. Math. **54** (1994), no. 2–3, 117–149.
10. Mohammad Hosseini Dolama and Éric Sopena, *On the oriented chromatic number of graphs with given excess*, Discrete Math. **306** (2006), no. 13, 1342–1350.
11. Mohammad Hosseini Dolama and Éric Sopena, *On the oriented chromatic number of Halin graphs*, Inform. Process. Lett. **98** (2006), no. 6, 247–252.
12. Keith Edwards, *The harmonious chromatic number and the achromatic number*, London Math. Soc. Lecture Note Ser., vol. 241, pp. 13–47, Cambridge Univ. Press, 1997.
13. ———, *The harmonious chromatic number of bounded degree graphs*, J. London Math. Soc. (2) **55** (1997), no. 3, 435–447.
14. ———, *A new upper bound for the harmonious chromatic number*, J. Graph Theory **29** (1998), no. 4, 257–261.
15. ———, *The harmonious chromatic number of complete r -ary trees*, Discrete Math. **203** (1999), no. 1–3, 83–99.
16. Paul Erdős, Alfréd Rényi, and Vera T. Sós, *On a problem of graph theory*, Studia Sci. Math. Hungar. **1** (1966), 215–235.
17. Louis Esperet and Pascal Ochem, *Oriented colorings of 2-outerplanar graphs*, Inform. Process. Lett. **101** (2007), no. 5, 215–219.
18. Hazel Everett and Arvind Gupta, *Acyclic directed hypercubes may have exponential diameter*, Inform. Process. Lett. **32** (1989), no. 5, 243–245.
19. Guillaume Fertin, André Raspaud, and Arup Roychowdhury, *On the oriented chromatic number of grids*, Inform. Process. Lett. **85** (2003), no. 5, 261–266.
20. Pierre Fraigniaud, Jean-Claude König, and Emmanuel Lazard, *Oriented hypercubes*, Networks **39** (2002), no. 2, 98–106.
21. Zoltán Füredi, Peter Horak, Chandra M. Pareek, and Xuding Zhu, *Minimal oriented graphs of diameter 2*, Graphs Combin. **14** (1998), no. 4, 345–350.
22. Johannes H. Hattingh, Michael A. Henning, and Elna Ungerer, *Graphs with small upper line-distinguishing and upper harmonious chromatic numbers*, J. Combin. Math. Combin. Comput. **47** (2003), 165–181.
23. ———, *Upper line-distinguishing and upper harmonious chromatic numbers of cycles*, J. Combin. Math. Combin. Comput. **45** (2003), 137–144.
24. Gyula O. H. Katona and Endre Szemerédi, *On a problem of graph theory*, Studia Sci. Math. Hungar. **2** (1967), 23–28.
25. William F. Klostermeyer and Gary MacGillivray, *Analogues of cliques for oriented coloring*, Discuss. Math. Graph Theory **24** (2004), no. 3, 373–387.
26. ———, *Homomorphisms and oriented colorings of equivalence classes of oriented graphs*, Discrete Math. **274** (2004), no. 1–3, 161–172.
27. ———, *Pushing vertices and oriented colorings*, Bull. Inst. Combin. Appl. **40** (2004), 49–58.

28. Alexandr V. Kostochka, Tomasz Luczak, Gábor Simonyi, and Eric Sopena, *On the minimum number of edges giving maximum oriented chromatic number*, DIMACS Ser. Discrete Math. Theoret. Comput. Sci., vol. 49, pp. 179–182, Amer. Math. Soc., 1999.
29. Alexandr V. Kostochka, Éric Sopena, and Xuding Zhu, *Acyclic and oriented chromatic numbers of graphs*, J. Graph Theory **24** (1997), no. 4, 331–340.
30. Marek Kubale, *Harmonious coloring of graphs*, Contemp. Math., vol. 352, pp. 95–104, Amer. Math. Soc., 2004.
31. Jiří Matoušek, *The number of unique-sink orientations of the hypercube*, Combinatorica **26** (2006), no. 1, 91–99.
32. Colin McDiarmid and Luo Xinhua, *Upper bounds for harmonious colorings*, J. Graph Theory **15** (1991), no. 6, 629–636.
33. Mirka Miller and Jozef Širáň, *Moore graphs and beyond: A survey of the degree/diameter problem*, Electron. J. Combin. (2005), DS14.
34. Jaroslav Nešetřil and André Raspaud, *Antisymmetric flows and strong colourings of oriented graphs*, Ann. Inst. Fourier (Grenoble) **49** (1999), no. 3, 1037–1056.
35. Jaroslav Nešetřil, André Raspaud, and Éric Sopena, *Colorings and girth of oriented planar graphs*, Discrete Math. **165-166** (1997), 519–530.
36. Jaroslav Nešetřil, Éric Sopena, and Laurence Vignal, *T-preserving homomorphisms of oriented graphs*, Comment. Math. Univ. Carolin. **38** (1997), no. 1, 125–136.
37. Pascal Ochem, *Oriented colorings of triangle-free planar graphs*, Inform. Process. Lett. **92** (2004), no. 2, 71–76.
38. André Raspaud and Éric Sopena, *Good and semi-strong colorings of oriented planar graphs*, Inform. Process. Lett. **51** (1994), no. 4, 171–174.
39. Attila Sali and Gábor Simonyi, *Oriented list colouring of undirected graphs*, DIMACS Ser. Discrete Math. Theoret. Comput. Sci., vol. 49, pp. 307–316, Amer. Math. Soc., 1999.
40. Robert Šámal, *Antisymmetric flows and strong oriented coloring of planar graphs*, Discrete Math. **273** (2003), no. 1–3, 203–209.
41. ———, *Antiflows, oriented and strong oriented colorings of graphs*, Arch. Math. (Brno) **40** (2004), no. 4, 335–343.
42. Éric Sopena, *The chromatic number of oriented graphs*, J. Graph Theory **25** (1997), no. 3, 191–205.
43. ———, *Oriented graph coloring*, Discrete Math. **229** (2001), no. 1–3, 359–369.
44. ———, *There exist oriented planar graphs with oriented chromatic number at least sixteen*, Inform. Process. Lett. **81** (2002), no. 6, 309–312.
45. Andrzej Szepietowski and Monika Targan, *A note on the oriented chromatic number of grids*, Inform. Process. Lett. **92** (2004), no. 2, 65–70.
46. K. Thilagavathi and Vernold Vivin. J, *Harmonious colouring of graphs*, Far East J. Math. Sci. **20** (2006), no. 2, 189–197.
47. David R. Wood, *Acyclic, star and oriented colourings of graph subdivisions*, Discrete Math. Theor. Comput. Sci. **7** (2005), no. 1, 37–50.

DEPARTAMENT DE MATEMÀTICA APLICADA II,
 UNIVERSITAT POLITÈCNICA DE CATALUNYA,
 BARCELONA, SPAIN
E-mail address: david.wood@upc.es